



High quality sustainable ingredients derived from Dried Black Soldier Fly Larvae (*Hermetia illucens*)

NUTRIENT	ENVIROBUG™ (Dry Larvae)*	ENVIROMEAL**	ENVIROOIL^
Moisture, %	8.00 (10.0 Max)	5.00 (10.0 Max)	-
Crude protein, %	35.0 (34.0 Min)	42 (40.0 Min)	-
Crude fiber, %	6.00 (10.0 Max)	6.50 (8.0 Max)	-
Acid detergent fiber, %	7.00	10.0	
Crude Fat, %	30.0 (26.0 Min)	Range: 12 – 20 (12.0 Min, 20.0 Max)	98.0 (98.0 Min)
Ash, %	7.00 (9.0 Max)	9.00 (12.0 Max)	-
Ca : P	2.14 : 1	2.09 : 1	-
Calcium, %	1.67	2.20	-
Phosphorus, %	0.78	1.05	-

Values are on *as-is basis* and are typical, based upon current averaged analysis;
Guaranteed Analyses are shown in **(bold)**.

Next update, Oct 2019

* Dried Black Soldier Fly larvae (whole) are defined for use in poultry, salmonid fish and wild bird feed as a source of protein and fat - T60.117(B) Dried Black Soldier Fly Larvae. AAFCO 2019 Official Publication. Association of American Feed Control Officials. Oxford, IN.

** Dried Black Soldier Fly larvae (ground) has been defatted and is defined for use in poultry and salmonid fish as a source of protein and fat - T60.117(B) Dried Black Soldier Fly Larvae. AAFCO 2019 Official Publication. Association of American Feed Control Officials. Oxford, IN.
Meal Crude Fat level will range based on extraction method used.

^ Dried Black Soldier Fly larvae oil is at this time not defined by the Association of American Feed Control Officials (AAFCO) as an ingredient for animal feeds.

EnviroFlight® products are all natural, free of added hormones and raised without antibiotics or pesticides.

FEEDING DIRECTIONS

This product is NOT a complete feed.

Consult your nutritionist for diet formulation to the specific needs of the intended species.

SHELF LIFE

Store product in dry area and maintain cool environmental conditions. After opening, ensure proper sealing of packaging to avoid moisture and air.

Use within 12 months of manufacturing (see Best Use by Date).

